

# Immunostimulatory Activity of *Hibiscus syriacus* L. Leaves in Mouse Macrophages, RAW264.7 cells, and Immunosuppressed Mice

Na Gyeong Geum<sup>1,3</sup>, Ju Hyeong Yu<sup>1</sup>, So Jung Park<sup>1</sup>, Min Yeong Choi<sup>1</sup>, Jae Won Lee<sup>2</sup>,  
Gwang Hun Park<sup>3</sup>, Hae-Yun Kwon<sup>4</sup> and Jin Boo Jeong<sup>5\*</sup>

<sup>1</sup>Graduate Student, <sup>2</sup>Post-doc and <sup>5</sup>Professor, Department of Medicinal Plant Resources,  
Andong National University, Andong 36729, Korea

<sup>3</sup>Researcher, Forest Medicinal Resources Research Center, National Institute of Forest Science, Yeongju 36040, Korea

<sup>4</sup>Researcher, Special Forest Resources Division, National Institute of Forest Science, Suwon 16631, Korea

**Abstract** - Under the COVID-19 pandemic, interest in immune enhancement is increasing. Although the immune-enhancing activity of plants of the genus *Hibiscus* has been reported, there is no study on the immune-enhancing activity of *H. syriacus*. Thus, in this study, we investigated the immune-enhancing activity of *Hibiscus syriacus* leaves (HSL) in mouse macrophages, RAW264.7 cells, and immunosuppressed mice. HSL increased the production of immunostimulatory factors such as nitric oxide (NO), inducible nitric oxide synthase (iNOS), interleukin-1 $\beta$  (IL-1 $\beta$ ), and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and activated the phagocytosis in RAW264.7 cells. The HSL-mediated production of immunostimulatory factors was dependent on toll-like receptor 4 (TLR4), p38, and c-Jun N-terminal kinase (JNK) in RAW264.7 cells. In the immunosuppressed mouse model, HSL increased the spleen index, the levels of the cytokines, and the numbers of lymphocytes, neutrophils, and monocytes. Taken together, HSL may be considered to have immune-enhancing activity and be expected to be used as a potential immune-enhancing agent.

**Key words** – *Hibiscus syriacus*, Immune enhancement, Macrophages

## Introduction

It is known that the immune response plays an important role in preventing and treating diseases caused by foreign pathogens (Hong *et al.*, 2017). Since immunostimulants can enhance the host's defense response to improve resistance to diseases caused by external pathogens, interest in the search for active substances that stimulate immune responses is increasing (Yu *et al.*, 2016).

Among the various immune cells, macrophages, which are innate immune cells, play an important role in the first defense against foreign pathogens (Geng *et al.*, 2018). Activated macrophages directly remove invading foreign pathogens through phagocytosis (Geng *et al.*, 2018). In addition, macrophages recognize foreign pathogens and provide antigen-specific information to T cells or B cells which are adaptive

immune cells (Zhou *et al.*, 2021). In addition, cytokines secreted by macrophages during immune response induce activation of adaptive immune cells such as T cells and B cells (Zhou *et al.*, 2021).

*Hibiscus syriacus* L. (*H. syriacus*) belonging to the Malvaceae family is the national flower of Republic of Korea and a medicinal plant that has been used to treat various diseases such as tinea, eczema, scabies, and dysentery (Shi *et al.*, 2014). *H. syriacus* has been reported to have various pharmacological activities such as anti-cancer activity, anti-inflammatory activity, antioxidant stress activity, and anti-brain disease (Karunaratne *et al.*, 2020). Although the immune-enhancing activity of plants of the genus *Hibiscus* has been reported (Geum *et al.*, 2021; Shen *et al.*, 2017), there is no study on the immune-enhancing activity of *H. syriacus*. Thus, we evaluated the immunostimulatory activity of *H. syriacus* and investigated its mechanism of action in mouse macrophages, RAW264.7 cells, and immunosuppressed mice.

\*Corresponding author. E-mail : jjb0403@anu.ac.kr  
Tel. +82-54-820-7757

## Materials and methods

### Chemical reagents

Specific inhibitors such as SB203580 (p38 inhibitor), PD98059 (ERK1/2 inhibitor), SP600125 (JNK inhibitor), C29 (TLR2 inhibitor) and TAK-242 (TLR4 inhibitor) were purchased from Sigma Aldrich (St. Louis, MO, USA). Neutral red for phagocytosis analysis of RAW264.7 macrophages and cyclophosphamide as immunosuppressant were purchased from Sigma Aldrich. Primary and secondary antibodies for Western blot analysis were purchased from Cell Signaling (Beverly, MA, USA).

### Sample preparation

The flowers, buds, leaves, and branches from *H. syriacus* were provided from Forest Medicinal Resources Research Center, Korea. Twenty grams of the flowers, flower buds, leaves, and branches from *H. syriacus* were immersed in 400 ml of distilled water and extracted for 12 h at 40°C. After extraction, the extracts were centrifuged for 10 min at 15,000 rpm, and then the supernatant was taken and lyophilized. The water extracts of flowers (HSF, yield=16.4%), flower buds (HSFB, yield=15.9%), leaves (HSL, yield=17.5%), and branches (HSB) from *H. syriacus* were stored in a refrigerator (-80 °C) until use. Each extract was dissolved in distilled water and treated to RAW264.7 cells and immunosuppressed mice.

### Cell culture

RAW264.7 cells purchased from American Type Culture Collection (Manassas, VA, USA) were maintained using Dulbecco's Modified Eagle medium (DMEM)/F-12 1:1 Modified medium (Lonza, Walkersville, MD, USA) containing 10% fetal bovine serum, 100 U/mL penicillin and 100 µg/mL streptomycin at 37°C under a humidified atmosphere of 5% CO<sub>2</sub>.

### MTT assay

To evaluate the cytotoxicity of HSL, an MTT assay was performed. Briefly, RAW264.7 cells ( $5 \times 10^5$  cells/mL) cultured for 24 h in a 96-well plate were treated with HSL, and then further cultured in an incubator (5% CO<sub>2</sub>, 37°C, Humidified atmosphere) for 24 h. After 24 h, 50 µl of MTT solution (1 mg

/mL, Sigma-Aldrich, St. Louis, MO, USA) was added to each well and then incubated in an incubator (5% CO<sub>2</sub>, 37°C, Humidified atmosphere) for 4 h. After 4 h, the culture medium was removed, and then 100 µl of dimethyl sulfoxide was added to each well to dissolve the purple formazan. The absorbance of dissolved purple formazan was measured at 570 nm using a UV/Visible spectrophotometer (Human Cop., Xma-3000PC, Seoul, Korea).

### Measurement of nitric oxide (NO)

The effect of HSF, HSFB, HSL, or HSB on the production of NO in RAW264.7 cells was evaluated using the Griess assay. Briefly, RAW264.7 cells were treated with HSF, HSFB, HSL, or HSB for 24 h. After 24 h, the cell culture supernatants and Griess reagent (Sigma Aldrich) were mixed and reacted at room temperature for 15 min. After 15 min, the absorbance was measured at 540 nm using a UV/Visible spectrophotometer (Human Cop., Xma-3000PC, Seoul, Korea).

### Measurement of phagocytic activity

The effect of HSL on the phagocytic activity in RAW264.7 cells was evaluated using a neutral red uptake assay. Briefly, RAW264.7 cells were treated with HSL for 24 h. After 24 h, RAW264.7 cells were washed three times with 1 X PBS and then stained with 0.01% neutral red solution for 2 h. After 2 h, RAW264.7 cells were washed three times with 1 X PBS and then absorbed neutral red was eluted with a lysis buffer (50% ethanol acid: 1% acetic acid = 1:1). The absorbance was measured at 540 nm using a UV/Visible spectrophotometer (Human Cop., Xma-3000PC, Seoul, Korea).

### Reverse transcription-polymerase chain reaction (RT-PCR)

The effect of HSF, HSFB, HSL, or HSB on the expression of inducible nitric oxide synthase (iNOS), interleukin-1β (IL-1β), and tumor necrosis factor-α (TNF-α) in RAW264.7 cells was evaluated using RT-PCR. Briefly, RAW264.7 cells were treated with HSF, HSFB, HSL, or HSB for 24 h. After 24 h, RAW264.7 cells were washed three times with 1 X PBS. Then, cDNA was synthesized using a Verso cDNA Kit (Thermo Scientific, Pittsburgh, PA, USA) from 1 µg of total RNA extracted from RAW264.7 cells using an RNeasy Mini Kit (Qiagen, Valencia, CA, USA). Then, PCR was performed

Table 1. The sequences of the primers used in this study

Primers	Sequence
iNOS	Forward 5'-ttgtcatcgcacttaggctggaa-3'
	Reverse 5'-gaccttcgcattagcatggaagc-3'
IL-1 $\beta$	Forward 5'-ggcaggcagtagtactcatt-3'
	Reverse 5'-cccaaggccacaggtattt-3'
TNF- $\alpha$	Forward 5'-tggaactggcagaagaggca-3'
	Reverse 5'-tgctcctccacttggtggtt-3'
GAPDH	Forward 5'-ggactgtggtcatgagccctcca-3'
	Reverse 5'-actcacggcaattcaacggcac-3'

using a PCR Master Mix Kit (Promega, Madison, WI, USA), and the sequences of the primers such as iNOS, IL-1 $\beta$ , TNF- $\alpha$ , and GAPDH were as Table 1. The density of mRNA bands was calculated using the software UN-SCAN-IT gel version 5.1 (Silk Scientific Inc. Orem, UT, USA).

#### Animals and experimental design

All animal procedures were conducted in accordance with the guidelines for the care and use of laboratory animals issued by Andong National University. Six-week-old male BALB/c mice were purchased from Orient (Sungnam, Korea). All mice were housed at  $25 \pm 2^\circ\text{C}$  and  $50 \pm 10\%$  humidity under a 12-h light/dark cycle. The mice were randomly assigned to 10 mice per group (Control group, Cyclophosphamide group, HSL 50 mg/kg group, HSL 100 mg/kg group, and HSL 200 mg/kg group). Cyclophosphamide (80 mg/kg mice) was intraperitoneally injected to mice daily for 3 days to suppress the immunity of mice. After 3 days, HSL was orally administered every day for 10 days. At the end of the experiment, mice were sacrificed, and then the spleens and cardiac blood from all mice were collected. The spleens were weighted immediately after sacrifice and the whole blood samples were placed in Vacutainer<sup>TM</sup> tubes containing EDTA (BD science, San Jose, NJ, USA). The numbers of white blood cells, neutrophils, lymphocytes, and monocytes were analyzed using a BC-5000 Vet Auto hematology analyzer (Nanshan, Shenzhen, China) in accordance with the manufacturer's recommendation. In addition, the levels of IL-1 $\beta$ , IL-6, and TNF- $\alpha$  from the serum of mice were analyzed by ELISA kit (Cayman Chemical, Ann Arbor, Michigan, USA) in accordance with manufacturer's protocols.

#### Statistical analysis

All the data are shown as mean  $\pm$  SD (standard deviation). Statistical significance was determined by Student's t-test (EXCEL, Microsoft Corporation, Redmond, Washington, USA). Differences with \* $P < 0.05$  or # $P < 0.05$  were considered statistically significant.

## Results and Discussion

#### HSL induces macrophage activation in RAW264.7 cells

It is reported that macrophages involved in the innate immune response recognize invaded foreign pathogens and remove them through phagocytosis (Kasimu *et al.*, 2017; Yu *et al.*, 2021). When foreign pathogens invade, macrophages secrete various immunostimulatory factors such as NO, iNOS, IL-1 $\beta$ , and TNF- $\alpha$  to eliminate them (Ma *et al.*, 2016; Siveen and Kuttan, 2009). It has been reported that NO produced by iNOS directly attacks foreign pathogens (Ren *et al.*, 2017). In addition, IL-1 $\beta$  and TNF- $\alpha$  have been known to strengthen the adaptive immune response by activating T cells and B cells (Zha *et al.*, 2018). Thus, it is reported that the activation of macrophages to induce secretion of these immunostimulatory factors contributes to host defense against external pathogens (Zhang *et al.*, 2013).

In this study, we investigated the effect of *H. syriacus* on the secretion of immunostimulatory factors such as NO, iNOS, IL-1 $\beta$ , and TNF- $\alpha$  in RAW264.7 cells to confirm whether *H. syriacus* has an immune-enhancing activity. First, we compared the effect of flowers (HSF), flower buds (HSFB), leaves (HSL), and branches (HSB) of *H. syriacus* L. on the secretion of the immunostimulatory factors. As shown in Fig. 1A, HSL and HSB dramatically increased the production of NO, iNOS, IL-1 $\beta$ , and TNF- $\alpha$  in RAW264.7 cells. HSF has been shown to increase the production of iNOS and TNF- $\alpha$  in RAW264.7 cells. However, HSFB did not increase any of NO, iNOS, IL-1 $\beta$ , and TNF in RAW264.7 cells. From the results (Fig. 1A), we selected HSL for further study. HSL dose-dependently increased the production of NO, iNOS, IL-1 $\beta$ , and TNF- $\alpha$  in RAW264.7 cells (Fig. 1B).

Since macrophages remove foreign pathogens through phagocytosis, phagocytosis activation of macrophages is considered to mean immune enhancement (We *et al.*, 2006).




Fig. 1. Effect of HSL on the macrophage activation in RAW264.7 cells. (A) After RAW264.7 cells were treated with HSF, HSFB, HSL or HSB for 24 h, the NO level and mRNA level were measured by Griess assay and RT-PCR, respectively. \*P < 0.05 compared to the cells without the treatment. (B) After RAW264.7 cells were treated with HSL for 24 h, the NO level and mRNA level were measured by Griess assay and RT-PCR, respectively. \*P < 0.05 compared to the cells without the treatment. (C) RAW264.7 cells were treated with HSL for 24 h. the absorbance of neutral red was investigated using a microplate reader. \*P < 0.05 compared to the cells without the treatment. (D) RAW264.7 cells were treated with HSL for 24 h. The cytotoxicity of HSL to RAW264.7 cells was performed using MTT assay. \*P < 0.05 compared to the cells without the treatment.

Thus, we investigated the effect of HSL on macrophage phagocytic activity in RAW264.7 cells. As shown in Fig. 1C, HSL dose-dependently increased the phagocytic activity in RAW264.7 cells. In addition, HSL did not affect the cytotoxicity of RAW264.7 cells (Fig. 1D). These results (Fig. 1A-1C) suggest that HSL may activate macrophages which may contribute to the enhancement of innate immune responses.

**HSL depends on TLR4 to increase the secretion of the immunostimulatory factors in RAW264.7 cells**

Toll-like receptors (TLRs) have been reported to play a key role in regulating macrophage activation (Swanson *et al.*, 2020). TLRs recognize pathogen-associated molecular patterns (PAMPs) of foreign pathogens and induce an inflammatory response to efficiently eliminate foreign pathogens (Moynagh, 2005; Swanson *et al.*, 2020). Thus, natural products that

stimulate TLRs were searched for macrophage activation, and it was reported that various natural products induce macrophage activation through stimulation of TLR2 and TLR4 among TLRs (Bai *et al.*, 2019; Shen *et al.*, 2017; Yang *et al.*, 2019; Zhang *et al.*, 2013). Thus, we investigated whether TLR2 and TLR4 were involved in the production of the immunostimulatory factors by HSL in RAW264.7 cells. As shown in Fig. 2, NO produced by HSL was reduced in TLR2 inhibition by C29, but there was no effect on the production of iNOS, IL-1β, and TNF-α in RAW264.7 cells. However, inhibition of TLR4 by TAK-242 markedly reduced the production of NO, iNOS, IL-1β, and TNF-α.

From these results, it is thought that HSL may induce the activation of macrophages mainly through the stimulation of TLR4.




Fig. 2. Effect of TLR2/4 on the HSL-induced production of immunostimulatory factors in RAW264.7 cells. RAW264.7 cells were pretreated with C29 (TLR2 inhibitor, 100  $\mu$ M) or TAK-242 (TLR4 inhibitor, 5  $\mu$ M) for 2 h and co-treated with HSL (30  $\mu$ g/mL) for 24 h. The NO level and mRNA level were investigated using Griess assay and RT-PCR. \* $P < 0.05$  compared to the cells without the treatment. # $P < 0.05$  compared to the cells with HSL alone.

### HSL depends on p38 and JNK to increase the secretion of the immunostimulatory factors in RAW264.7 cells

Mitogen-activated protein kinases (MAPKs), composed of p38, extracellular signal-regulated protein kinase 1/2 (ERK1/2), and c-Jun N-terminal kinase (JNK), are known as one of the major signaling pathways involved in the secretion of immunostimulatory factors by macrophages (Shen *et al.*, 2017). Moreover, it has been reported that TLRs induce the activation of macrophages by activating MAPK signaling (Shen *et al.*, 2017). Thus, we investigated whether MAPKs were involved in the production of the immunostimulatory factors by HSL in RAW264.7 cells. As shown in Fig. 3, there was no change in the production of immunostimulatory factors such as NO, iNOS, IL-1 $\beta$ , and TNF- $\alpha$  in the inhibition of ERK1/2 by PD98059. However, inhibition of p38 by SB203580 decreased the production of NO and IL-1 $\beta$  by HSL, and inhibition of JNK by SP600125 decreased the production of NO, iNOS, IL-1 $\beta$ , and TNF- $\alpha$  by HSL.

From these results, it is thought that p38 and JNK are signaling pathways involved in the HSL-mediated activation of macrophages.

### HSL increases immune enhancement-related markers in immunosuppressed mouse models

To verify the immune-enhancing activity of HSL in an animal




Fig. 3. Effect of MAPK signaling pathway on the HSL-induced production of immunostimulatory factors in RAW264.7 cells. RAW264.7 cells were pretreated with PD98059 (ERK1/2 inhibitor, 40  $\mu$ M), SB203580 (p38 inhibitor, 40  $\mu$ M) or SP600125 (JNK inhibitor, 40  $\mu$ M) for 2 h and then co-treated with HSL (30  $\mu$ g/mL) for 24 h. NO level and mRNA level were measured by Griess assay and RT-PCR, respectively. \* $P < 0.05$  compared to the cells without the treatment. # $P < 0.05$  compared to the cells with HSL alone.

model, immunosuppressed mice were treated with HSL, and then the immune enhancement-related markers were analyzed.

The spleen is an organ in the body that plays an important role in the innate and acquired immune systems (Aliyu *et al.*, 2021). It is known that various immune cells such as B cells, natural killer cells, and macrophages exist in the spleen (Aliyu *et al.*, 2021). Therefore, the increase in the spleen index in the immunosuppressed mouse model has been utilized as an immune enhancement index. In this study, we found that treatment of cyclophosphamide (CP) as an immunosuppressant reduced the splenic index, but HSL restored the reduced splenic index by CP (Fig. 4A).

Blood is called the pipeline of the immune system because it contains various types of immune cells such as lymphocytes, neutrophils, and monocytes and immunostimulatory factors such as IL-1 $\beta$ , IL-6, and TNF- $\alpha$  (Geum *et al.*, 2021). Thus, we analyzed the number of lymphocytes, neutrophils, and monocytes in the blood. As a result (Fig. 4B), treatment with CP alone decreased the number of lymphocytes, neutrophils, and monocytes compared to the normal group, but administration of HSL restored the reduced immune cells by CP such as lymphocytes, neutrophils, and monocytes. In addition, the administration of HSL increased the levels of IL-1 $\beta$ , IL-6, and TNF- $\alpha$  in the serum of the blood.




Fig. 4. Effect of HSL on the immune-enhancement-related markers in immunosuppressed mice. (A) The spleen index was calculated as spleen weight/body weight. (B) The numbers of lymphocytes, neutrophils, and monocytes were analyzed using a BC-5000 Vet Auto hematology analyzer. (C) The levels of IL-1 $\beta$ , IL-6 and TNF- $\alpha$  were measured from the serum of the blood using ELISA kits. \*P < 0.05 compared to the cells without the treatment. #P < 0.05 compared to the cells with HSL alone.

In summary, HSL increased the production of NO, iNOS, IL-1 $\beta$ , and TNF- $\alpha$  through the TLR4-dependent activation of p38 and JNK in RAW264.7 cells. In addition, HSL was shown to increase the spleen index, the numbers of lymphocytes, neutrophils, and monocytes, and the levels of the cytokines (IL-1 $\beta$ , IL-6 and TNF- $\alpha$ ) in the immunosuppressed mouse model. These findings indicate that HSL may have immunostimulatory activity. Therefore, HSL may be considered to have immune-enhancing activity, and HSL is expected to be used as a potential immune-enhancing agent.

### Acknowledgments

This work was supported by a grant from National Institute of

Forest Science in 2021 (project number: FG0403-2018-01-2021) and R&D Program for Forest Science Technology (Project No. 2021377C10-2123-BD02) provided by Korea Forest Service (Korea Forestry Promotion Institute).

### Conflicts of Interest

The authors declare that they have no conflict of interest.

### References

Aliyu, M., F. Zohora and A.A. Saboor-Yaraghi. 2021. Spleen in innate and adaptive immunity regulation. *AIMS Allergy Immunol.* 5(1):1-17.

- Bai, Y., Y. Jiang, T. Liu, F. Li, J. Zhang, Y. Luo, L. Zhang, G. Yan, Z. Feng, X. Li, X. Wang and W. Hu. 2019. Xinjiang herbal tea exerts immunomodulatory activity via TLR2/4-mediated MAPK signalling pathways in RAW264.7 cells and prevents cyclophosphamide-induced immunosuppression in mice. *J. Ethnopharmacol.* 228:179-187.
- Geng, L., W. Hu, Y. Liu, J. Wang and Q. Zhang. 2018. A heteropolysaccharide from *Saccharina japonica* with immunomodulatory effect on RAW264.7 cells. *Carbohydr. Polym.* 201:557-565.
- Geum, N.G., J.H. Yu, J.H. Yeo, M.Y. Choi, J.W. Lee, J.K. Baek and J.B. Jeong. 2021. Immunostimulatory activity and anti-obesity activity of *Hibiscus manihot* leaves in mouse macrophages, RAW264.7 cells and mouse adipocytes, 3T3-L1 cells. *J. Funct. Foods* 87:104803.
- Hong, S.H., J.M. Ku, H.I. Kim, C.W. Ahn, S.H. Park, H.S. Seo, Y.C. Shin and S.G. Ko. 2017. The immune-enhancing activity of *Cervus nippon manchuricus* extract (NGE) in RAW264.7 macrophage cells and immunosuppressed mice. *Food Res. Int.* 99(1):623-629.
- Karunaratne, W.A.H.M., K.T. Lee, Y.H. Choi, C.Y. Jin and G.Y. Kim. 2020. Anthocyanins isolated from *Hibiscus syriacus* L. attenuate lipopolysaccharide-induced inflammation and endotoxic shock by inhibiting the TLR4/MD2-mediated NF- $\kappa$ B signaling pathway. *Phytochemistry* 76: 153237.
- Kasimu, R., C. Chen, X. Xie and X. Li. 2017. Water-soluble polysaccharide from *Erythronium sibiricum* bulb: structural characterisation and immunomodulating activity. *Int. J. Biol. Macromol.* 105:452-462.
- Ma, X.L., M. Meng, L.R. Han, D. Cheng, X.H. Cao and C.L. Wang. 2016. Structural characterization and immunomodulatory activity of *Grifola frondosa* polysaccharide via toll-like receptor 4-mitogen-activated protein kinases-nuclear factor  $\kappa$ B pathways. *Food Funct.* 7:2763-2772.
- Moynagh, P.N. 2005. TLR signalling and activation of IRFs: revisiting old friends from the NF-kappaB pathway. *Trends Immunol.* 26(9):469-479.
- Ren, Z., T. Qin, F. Qiu, T. Song, D. Lin, Y. Ma, J. Li and Y. Huang. 2017. Immunomodulatory effects of hydroxyethylated *Hericium erinaceus* polysaccharide on macrophages RAW 264.7. *Int. J. Biol. Macromol.* 105(1):879-885.
- Shen, C.Y., W.L. Zhang and J.G. Jiang. 2017. Immune-enhancing activity of polysaccharides from *Hibiscus sabdariffa* Linn. via MAPK and NF- $\kappa$ B signaling pathways in RAW264.7 cells. *J. Funct. Foods* 34:118-129.
- Shi, L.S., C.H. Wu, T.C. Yang, C.W. Yao, H.C. Lin and W.L. Chang. 2014. Cytotoxic effect of triterpenoids from the root bark of *Hibiscus syriacus*. *Fitoterapia* 97:184-191.
- Siveen, K.S. and G. Kuttan. 2009. Role of macrophage in tumour progression. *Immunol. Lett.* 123:97-102.
- Swanson, L., G.D. Katkar, J. Tam, R.F. Pranadinata, Y. Chareddy, J. Coates, M.S. Anandachar, V. Castillo, J. Olson, V. Nizet, I. Kufareva, S. Das and P. Ghosh. 2020. TLR4 signaling and macrophage inflammatory responses are dampened by GIV/Girdin. *PNASU.* 117(43):26895-26906.
- We, M., T. Cheng, S. Cheng, T. Lian, L. Wang and S. Chiou. 2006. Immunomodulatory properties of *Grifola frondosa* in submerged culture. *J. Agri. Food Chem.* 54:2906-2914.
- Yang, F., X. Li, Y. Yang, S.M. Ayivi-Tosuh, F. Wang, H. Li and G. Wang. 2019. A polysaccharide isolated from the fruits of *Physalis alkekengi* L. induces RAW264.7 macrophages activation via TLR2 and TLR4-mediated MAPK and NF- $\kappa$ B signaling pathways. *Int. J. Biol. Macromol.* 140:895-906.
- Yu, J.H., N.G. Geum, J.H. Yeo and J.B. Jeong. 2021. Immune-enhancing and anti-obesity effect of *Abelmoschus manihot* root extracts. *Korean J. Plant Res.* 34(5):411-419.
- Yu, Z., M. Kong, P. Zhang, Q. Sun and K. Chen. 2016. Immune-enhancing activity of extracellular polysaccharides isolated from *Rhizopus nigricans*. *Carbohydr. Polym.* 148:318-325.
- Zha, Z., S.Y. Wang, W. Chu, Y. Lv, H. Kan, Q. Chen, L. Zhong, L. Yue, J. Xiao, Y. Wang and H. Yin. 2018. Isolation, purification, structural characterization and immunostimulatory activity of water-soluble polysaccharides from *Lepidium meyenii*. *Phytochemistry* 147:184-193.
- Zhang, S., S. Nie, D. Huang, J. Huang, Y. Wang and M. Xie. 2013. Polysaccharide from *Ganoderma atrum* evokes anti-tumor activity via toll-like receptor 4-mediated NF- $\kappa$ B and mitogen-activated protein kinase signaling pathways. *J. Agri. Food Chem.* 61:3676-3682.
- Zhou, R., L. Teng, Y. Zhu, C. Zhang, Y. Yang and Y. Chen. 2021. Preparation of *Amomum longiligulare* polysaccharides 1- PLGA nanoparticle and its immune enhancement ability on RAW264.7 cells. *Int. Immunopharmacol.* 99:108053.

(Received 31 May 2022 ; Revised 28 June 2022 ; Accepted 28 June 2022)